

California Pest Rating Profile for

Xyleborus monographus (Fabricius): Mediterranean oak borer

Coleoptera: Curculionidae

Previous Pest Rating: A

Pest Rating: B as of 02/05/2021

Comment Period: **12/22/2020 – 2/5/2021**

Initiating Event:

Xyleborus monographus was found infesting several valley oak (*Quercus lobata*) trees in Calistoga, Napa County, California in 2019. An A-rating was assigned to this species in March 2020. Since then, trapping and surveys of trees have expanded the known distribution of this beetle in California to four counties. The risk this species poses to California needs to be re-assessed. Therefore, a pest rating proposal is needed.

History & Status:

Background: Adult *X. monographus* range from 1.8 to 3.6 mm in length. Adults tunnel into trees that are often reported to be recently dead or stressed. Host trees reported worldwide include chestnut (including *Castanea sativa*), beech (including *Fagus orientalis*), and oak (including *Quercus douglassii*, *Q. frainetto*, *Q. kelloggii*, *Q. lobata*, *Q. pyrenaica*, *Q. rober*, *Q. rubra*, and *Q. suber*) species (California Department of Food and Agriculture; C. Ewing, pers. comm.; Lombardero, 1995; Sarikaya and Kavakli, 2018). Additional genera of deciduous trees are mentioned by Schedl (1964), but he states that oaks are the most important hosts. Adult females tunnel into trunks and branches, usually at points where the bark is thin or cracked. The entrance holes measure 1.3-1.5 mm in diameter (Schedl, 1964). Adult females carry with them fungal spores in structures called mycangia that are located (in this species) on their mandibles. These spores inoculate the branching tunnels made by the beetles.

Beetle larvae feed on the fungus “garden.” Besides being the primary food of the beetle larvae and adults, the fungi may be pathogenic to living trees. One of the fungal species associated with *X. monographus* in Europe and in the 2019 Calistoga infestation is *Raffaelea montetyi* (A. Eskalen, pers. comm.; Gebhardt et al., 2004). This fungus is reported to be pathogenic to cork oak and in experiments it killed inoculated seedlings within two months (Inácio et al., 2012). The trees associated with the 2019 finds of *X. monographus* in Calistoga showed symptoms including wilting, defoliation, and broken branches, and the infested wood was discolored, presumably by the fungus (A. Eskalen, pers. comm.). Infested oak trees in various locations in the infested area of California show crown dieback that appears to progress in a downward direction from outer branches (C. Ewing, pers. comm.). *Raffaelea montetyi* (but not *X. monographus*, although it was likely present but appears to have been misidentified as another scolytine species) was reported in 2018 in Calistoga infesting *Q. lobata* (California Forest Pest Council). This appears to be the first report of this fungus in North America and the fungus is presumed to have been introduced at the same time as the beetle.

In addition to injuring or killing living trees, *X. monographus* is reported to damage cut oak wood (Franjević et al., 2016).

Xyleborus species have sibling mating. Therefore, females can mate with their brothers prior to leaving the gallery (with fungus in their mycangia) and flying to a new tree to initiate gallery construction. Males have reduced wings, cannot fly, and are not involved with the construction of new galleries (Schedl, 1964). In addition, *Xyleborus* species are parthenogenic. An unmated female can lay unfertilized eggs that develop into males that the female can then mate with to produce fertilized eggs (Gohli et al., 2016).

Worldwide Distribution: *Xyleborus monographus* is reported from Africa (Algeria and Morocco), Europe (including France, Italy, Norway, Slovenia, and Spain), Palearctic Asia (including Israel, Korea, and Turkey), and North America (only in the United States, and only in California) (Buse et al., 2013; Choo and Woo, 1985; Freeman and Grancher, 2014; Hardersen et al., 2014; Hauptman et al., 2019; Lombardero, 1995; Olberg, 2007; Rabaglia et al., 2020; Sarikaya and Kavakli, 2018). The Calistoga,

California finds in 2019 appear to be the first known establishment of the species in the United States. One specimen was trapped in Portland, Oregon in 2018, but no further specimens appear to have been found there, suggesting this find may not have represented an infestation (Oregon Department of Agriculture). Molecular data (Cytochrome oxidase I gene) suggests France may be the origin of the California infestation (C. Ewing, pers. comm.).

Official Control: : *Xyleborus monographus* is listed as a quarantine pest by Japan and it is considered reportable by the United States Department of Agriculture (Food and Agriculture Organization of the United Nations; USDA-APHIS).

California Distribution: *Xyleborus monographus* has been found infesting trees and has been caught in ethanol traps in Lake, Napa, Sacramento, and Sonoma counties (California Department of Food and Agriculture). Based on the find of its ambrosia fungus in 2018 in Calistoga, it appears to have been established in California for at least several years and observations of old wood with characteristic galleries suggests this beetle has been in California longer than that.

California Interceptions: *Xyleborus monographus* was trapped near a wood processing site in Richmond (Contra Costa County) in 2018 (California Department of Food and Agriculture).

The risk *Xyleborus monographus* poses to California is evaluated below.

Consequences of Introduction:

- 1) **Climate/Host Interaction:** Although *X. monographus* feeds on its symbiotic fungi, it attacks a variety of oaks and other trees in the family Fagaceae. The most commonly-attacked tree in California is valley oak (*Quercus lobata*), which is widespread in this state. *Xyleborus monographus* is found in a variety of climates, including Mediterranean. It is likely capable of establishing over much of California. Therefore, it receives a **High (3)** in this category.

- Low (1) Not likely to establish in California; or likely to establish in very limited areas.
- Medium (2) may be able to establish in a larger but limited part of California.
- **High (3) likely to establish a widespread distribution in California.**

2) **Known Pest Host Range:** *Xyleborus monographus* feeds on symbiotic fungi, but for the purpose of this proposal, the trees it attacks and establishes galleries in will be considered hosts. These trees include several genera in the family Fagaceae (but see Uncertainty, below). Therefore, it receives a **Medium (2)** in this category.

- Low (1) has a very limited host range.
- **Medium (2) has a moderate host range.**
- High (3) has a wide host range.

3) **Pest Reproductive and Dispersal Potential:** Like others in the genus, female *X. monographus* are presumed to mate with their siblings and be parthenogenetic. A single mated or unmated female is therefore capable of establishing a new infestation. Ambrosia beetles can be moved in infested wood and the females can fly. Therefore, it receives a **High (3)** in this category.

- Low (1) does not have high reproductive or dispersal potential.
- Medium (2) has either high reproductive or dispersal potential.
- **High (3) has both high reproduction and dispersal potential.**

4) **Economic Impact.** *Xyleborus monographus* has a preference for older trees and mainly attacks branches with a diameter of at least seven inches, although branches as narrow as two inches are attacked (C. Ewing, pers. comm.). Therefore, this beetle does not appear to pose a significant risk to tree nurseries. As a reportable pest, it may trigger quarantines. Therefore, it receives a **Low (1)** in this category.

Economic Impact: C

- A. The pest could lower crop yield.
- B. The pest could lower crop value (includes increasing crop production costs).
- C. The pest could trigger the loss of markets (includes quarantines).**
- D. The pest could negatively change normal cultural practices.
- E. The pest can vector, or is vectored, by another pestiferous organism.
- F. The organism is injurious or poisonous to agriculturally important animals.
- G. The organism can interfere with the delivery or supply of water for agricultural uses.

Economic Impact Score: Low

– **Low (1) causes 0 or 1 of these impacts.**

– Medium (2) causes 2 of these impacts.

– High (3) causes 3 or more of these impacts.

- 5) **Environmental Impact.** *Xyleborus monographus* and its fungal symbiont apparently attack living oak trees. In California, this beetle-fungus mutualism will be exposed to species of oaks that it has not previously been exposed to and damage to some of these species could be more severe than that observed in the beetle's native range. Oaks are important components of forests and woodlands in California and valley oak (*Q. lobata*), which is already known to be attacked, is widely distributed in California and an important component of oak woodlands. Some species of oaks in this state are rare. Oaks (including *Q. lobata*) provide habitat for threatened species (Beckman and Jerome, 2017). Oak trees in California, including *Q. lobata*, have historically been used by Native Americans for food and other purposes. Oaks, especially the large heritage oaks, are important landscape trees and add to the value of property. Therefore, *X. monographus* receives a **High (3)** in this category.

Environmental Impact: A, B, C, E

- A. The pest could have a significant environmental impact such as lowering biodiversity, disrupting natural communities, or changing ecosystem processes.**
- B. The pest could directly affect threatened or endangered species.**
- C. The pest could impact threatened or endangered species by disrupting critical habitats.**
- D. The pest could trigger additional official or private treatment programs.
- E. The pest significantly impacts cultural practices, home/urban gardening or ornamental plantings.**

Environmental Impact Score: High (3)

- Low (1) causes none of the above to occur.
- Medium (2) causes one of the above to occur.
- **High (3) causes two or more of the above to occur.**

Consequences of Introduction to California for *Xyleborus monographus*: Medium (12)

Add up the total score and include it here.

- Low = 5-8 points
- Medium = 9-12 points**
- High = 13-15 points

6) Post Entry Distribution and Survey Information: *Xyleborus monographus* is established in Lake, Napa, Sacramento, and Sonoma counties. These areas represent the Coast Range mountains and the Central Valley of California. It receives a **Medium (-2)** in this category.

- Not established (0) Pest never detected in California, or known only from incursions.
- Low (-1) Pest has a localized distribution in California, or is established in one suitable climate/host area (region).

–**Medium (-2) Pest is widespread in California but not fully established in the endangered area, or pest established in two contiguous suitable climate/host areas.**

–High (-3) Pest has fully established in the endangered area, or pest is reported in more than two contiguous or non-contiguous suitable climate/host areas.

Final Score:

7) The final score is the consequences of introduction score minus the post entry distribution and survey information score: Medium (10)

Uncertainty:

It is possible that *X. monographus* is more widely distributed in California than is currently known. Traps for exotic wood boring beetles are used in certain California counties as part of a CAPS (Cooperative Agricultural Pest Survey) project. Although *X. monographus* is not a specific target of the California survey, the traps used include Lindgren funnels with ethanol lures, which attract *X. monographus* (California Department of Food and Agriculture; Galko et al., 2014). Over the past two years, these traps have been placed in Alameda, Contra Costa, Monterey, San Diego, San Francisco, San Joaquin, San Luis Obispo, San Mateo, Santa Barbara, Santa Clara, and Santa Cruz counties. The trap catches are reviewed by qualified personnel who would likely recognize exotic species of ambrosia beetles. There is uncertainty regarding the presence of *X. monographus* in California counties that are not routinely trapped for wood boring beetles, especially around the area currently known to be infested (Lake, Napa, Sonoma, and Sacramento counties). There is less uncertainty regarding its presence in the counties included in the survey, including the San Francisco Bay area, where 16 sites were trapped with ethanol lures in 2018-2019 (C. Takahashi, pers. comm.).

Remnant oak trees serve as habitat for bats (and likely other animals) in California vineyards (Polyakov et al., 2019). Infestation and death of such trees could therefore impact animals associated with these oak trees, which could in turn impact ecosystem services, such as pest control, provided by these animals. The host range of *Xyleborus monographus* could be much narrower or broader in California than has been observed in Europe. If so, its impact in the state could be over- or underestimated in this proposal. It may not be a competent attacker of oak trees other than the

species already found to be infested here and in Europe. However, the valley oak, the most commonly-reported host of this beetle in California, is a widespread and important species.

Conclusion and Rating Justification:

Xyleborus monographus is an ambrosia beetle that is known to be established in four counties in California. It vectors a pathogenic fungus that is capable of killing oak trees and it has most often been found on valley oak, a widespread tree in California. Its ultimate impact on the state's trees is not yet known. However, it is already widespread in multiple counties in different ecological regions and eradication does not appear to be feasible at this point. For these reasons, a "B" rating is justified.

References:

Beckman, E., and Jerome, D. 2017. *Quercus lobata*. The IUCN Red List of Threatened Species.

Accessed December 10, 2019:

<https://www.iucnredlist.org/species/61983021/61983023>

Buse, J., Assmann, T., Friedman, A. -L. -L., Rittner, O., and Pavlicek, T. 2013. Wood-inhabiting beetles (Coleoptera) associated with oaks in a global biodiversity hotspot: a case study and checklist for Israel. *Insect Conservation and Diversity* 6:687-703.

California Department of Food and Agriculture. Pest and damage record database. Accessed December 7, 2020:

<https://pdr.cdfa.ca.gov/PDR/pdrmainmenu.aspx>

California Forest Pest Council. 2018 California forest pest conditions. Accessed November 22, 2019:

<https://bof.fire.ca.gov/media/9196/full-12-2018-pest-conditions-report-ada.pdf>

Choo, H. Y. and Woo, K. S. 1985. A list of Korean bark and ambrosia beetles, and their host plants. *Korean Journal of Plant Protection* 24:163-167.

Food and Agriculture Organization of the United Nations. List of regulated pests from Japan.

Accessed November 27, 2019:

<https://www.ippc.int/en/countries/japan/reportingobligation/3>

Franjević, M., Poršinsky, T., and Đuka, A. 2016. Integrated oak timber protection from ambrosia bark beetles: economic and ecological importance in harvesting operations. *Croatian Journal of Forest Engineering* 37:353-364.

Freeman, J. -C. and Grancher, C. 2014. La forêt domaniale de Bastard (Pyrénées-Atlantiques) : un espace riche en Coléoptères saproxyliques aux portes de Pau. Bulletin de la Société Linnéenne de Bordeaux 42:77-88.

Galko, J., Nikolov, C., Kimoto, T., Kunca, A., Gubka, A., Vahula, J., Zúbrik, M., and Ostrihoň, M. 2014. Attraction of ambrosia beetles to ethanol baited traps in a Slovakian oak forest. Biologia 69:1376-1383.

Gebhardt, H., Begerow, D., and Oberwinkler, F. 2004. Identification of the ambrosia fungus of *Xyleborus monographus* and *X. dryographus* (Coleoptera: Curculionidae, Scolytinae). Mycological Progress 3:95-102.

Gohli, J., Selvarajah, T., Kirkendall, L. R., and Jordal, B. J. 2016. Globally distributed *Xyleborus* species reveal current intercontinental dispersal in a landscape of ancient worldwide distributions. BMC Evolutionary Biology 16:1-12.

Hardersen, S., Curletti, G., Leseigneur, L., Platia, G., Liberti, G., Leo, P., Cornacchia, P., and Gatti, E. 2014. Spatio-temporal analysis of beetles from the canopy and ground layer in an Italian lowland forest. Bulletin of Insectology 67:87-97.

Hauptman, T., Pavlin, R., Grošelj, P., and Jurc, M. 2019. Distribution and abundance of the alien *Xylosandrus germanus* and other ambrosia beetles (Coleoptera: Curculionidae, Scolytinae) in different forest stands in central Slovenia. iForest 12:451-458.

Inácio, M. L., Henriques, J., Lima, A., and Sousa, E. 2012. Ophiostomatoid fungi associated with cork oak mortality in Portugal. IOBC/WPRS Bulletin 76:89-92.

Lombardero, M. J. 1995. Plantas huésped y escolítidos (Col.: Scolytidae) en Galicia (Noroeste de la Península Ibérica). Boletín de Sanidad Vegetal – Plagas 21:357-370.

Olberg, S. 2007. *Agilus cyanescens* Ratzburg, 1837 (Buprestidae) and *Xyleborus monographus* (Fabricius, 1792) (Curculionidae) – two new but probably extinct Norwegian Coleoptera. Norwegian Journal of Entomology 54:115-116.

Oregon Department of Agriculture. Annual report 2018. Accessed November 26, 2019: <https://www.oregon.gov/ODA/shared/Documents/Publications/NurseryChristmasTree/PlantProgrmsAnnualReport.pdf>

Polyakov, A. Y., Weller, T. J., and Tietje, W. D. 2019. Remnant trees increase bat activity and facilitate the use of vineyards by edge-space bats. Agriculture, Ecosystems and Environment 281:56-63.

Rabaglia, R. J., Smith, S. L., Rugman-Jones, P., Digirolomo, M. F., Ewing, C., and Eskalen, A. 2020. Establishment of a non-native xyleborine ambrosia beetle, *Xyleborus monographus* (Fabricius) (Coleoptera: Curculionidae: Scolytinae), new to North American in California. *Zootaxa* 4786:269-276.

Sarikaya, O. and Kavakli, S. A. 2018. Faunistic observations on Scolytinae (Coleoptera: Curculionidae) in Afyonkarahisar region of Turkey. *American Journal of Engineering Research* 7:277-282.

Schedl, W. Biologie des gehöckerten Eichenholzbohrers, *Xyleborus monographus* Fab. (Scolytidae, Coleoptera). *Zeitschrift für Angewandte Entomologie* 53:411–428.

USDA-APHIS. U.S. regulated plant pest table. Accessed December 9, 2020:
<https://www.aphis.usda.gov/aphis/ourfocus/planthealth/import-information/rppl/rppl-table>.

Responsible Party:

Kyle Beucke, 2800 Gateway Oaks, Suite #200, Sacramento, CA, 95833, 916-403-6741,
permits[@]cdfa.ca.gov

***Comment Period: 12/22/2020 – 2/5/2021**

***NOTE:**

You must be registered and logged in to post a comment. If you have registered and have not received the registration confirmation, please contact us at [permits\[@\]cdfa.ca.gov](mailto:permits[@]cdfa.ca.gov).

Comment Format:

- ❖ Comments should refer to the appropriate California Pest Rating Proposal Form subsection(s) being commented on, as shown below.

Example Comment:

Consequences of Introduction: 1. Climate/Host Interaction: [Your comment that relates to “Climate/Host Interaction” here.]

- ❖ Posted comments will not be able to be viewed immediately.

- ❖ Comments may not be posted if they:

Contain inappropriate language which is not germane to the pest rating proposal;

Contains defamatory, false, inaccurate, abusive, obscene, pornographic, sexually oriented, threatening, racially offensive, discriminatory or illegal material;



Violates agency regulations prohibiting sexual harassment or other forms of discrimination;

Violates agency regulations prohibiting workplace violence, including threats.

- ❖ Comments may be edited prior to posting to ensure they are entirely germane.
- ❖ Posted comments shall be those which have been approved in content and posted to the website to be viewed, not just submitted.

Pest Rating: B